



## EMBRACE 3.6L

### Wheat

For use on wheat to suppress *Fusarium* head blight (scab)

### SECTION 18 SPECIFIC EXEMPTION

File Symbol: 07-MI-11

### FOR DISTRIBUTION AND USE ONLY IN THE **STATE OF MICHIGAN** UNDER SECTION 18 SPECIFIC EXEMPTION

This labeling must be in possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions including precautions when spraying in the vicinity of aquatic areas, on the registered product label for EMBRACE 3.6L (EPA Reg No. 1381-203) and EMBRACE 3.6L (EPA Reg No. 42750-99-1381).

Any adverse effects resulting from the use of EMBRACE 3.6L under this specific exemption must be immediately reported to the State of Michigan Department of Agriculture.

#### EFFECTIVE PERIOD:

This labeling expires on June 25, 2007.

#### DIRECTIONS FOR USE

A maximum of 250,000 acres of wheat may be treated under this exemption.

#### RECOMMENDED APPLICATIONS

CROP	DISEASES	RATE OF EMBRACE 3.6L
Wheat (including durum)	<i>Fusarium</i> head blight (scab)	4 fl oz per acre

#### USE DIRECTIONS

Apply EMBRACE 3.6L as a preventive foliar spray. For optimum disease suppression, the lowest rate of a spray surfactant should be tank-mixed with EMBRACE 3.6L. EMBRACE 3.6L may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 3 gallons of spray solution per acre by aircraft spray equipment.

Application through any type of irrigation equipment is prohibited.

Do not make more than 1 application of a tebuconazole-containing product to wheat per season. One application may be made using ground or aerial equipment at a rate of 4 fluid ounces of formulated product (1.8 oz a.i.) per acre per year. EMBRACE 3.6L may be applied up through full head emergence (Feekes 10.5.1). Application may not be made within 30 days of harvest. Straw

cut after harvest may be fed or used for bedding. Do not enter treated field for 12 hours after application.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to freshwater, estuarine and marine fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

For emergencies call CHEMTREC toll free 24 Hours A Day 1-800-424-9300

For PRODUCT USE information, visit at [www.agriliance.com](http://www.agriliance.com)

As with any crop protection product, always read and follow label directions.

Registered by:  
Agriliance LLC  
PO Box 64089  
St. Paul, MN 55164